

# Augmented pressor and sympathetic responses to skeletal muscle metaboreflex activation in type 2 diabetes patients

Holwerda, Seth W; Restaino, Robert M; Manrique, Camila; Lastra, Guido; Fisher, James P; Fadel, Paul J

DOI:

[10.1152/ajpheart.00636.2015](https://doi.org/10.1152/ajpheart.00636.2015)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Holwerda, SW, Restaino, RM, Manrique, C, Lastra, G, Fisher, JP & Fadel, PJ 2016, 'Augmented pressor and sympathetic responses to skeletal muscle metaboreflex activation in type 2 diabetes patients', *American Journal of Physiology Heart and Circulatory Physiology*, vol. 310, no. 2, pp. H300-9.  
<https://doi.org/10.1152/ajpheart.00636.2015>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

Eligibility for repository: Checked on 10/3/2016

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

**Augmented pressor and sympathetic responses to skeletal muscle metaboreflex activation  
in type 2 diabetes patients**

Seth W. Holwerda<sup>1</sup>, Robert M. Restaino<sup>1</sup>, Camila Manrique<sup>2</sup>, Guido Lastra<sup>2</sup>, James P. Fisher<sup>4</sup>,  
Paul J. Fadel<sup>1,3</sup>

<sup>1</sup>*Department of Medical Pharmacology and Physiology*, <sup>2</sup>*Department of Medicine*, <sup>3</sup>*Dalton  
Cardiovascular Research Center, University of Missouri, Columbia, MO*, <sup>4</sup>*School of Sport,  
Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK*

Running Title: augmented metaboreflex activation in T2D

Keywords: exercise, hypertension, muscle sympathetic nerve activity, post exercise ischemia,  
blood pressure

Correspondence:

Paul J. Fadel, PhD

Department of Kinesiology

College of Nursing and Health Innovation

University of Texas at Arlington

500 W. Nedderman #115, Arlington, TX 76019

1-817-272-3288

fadelp@uta.edu

## ABSTRACT

Previous studies have reported exaggerated increases in arterial blood pressure during exercise in type 2 diabetes (T2D) patients. However, little is known regarding the underlying neural mechanism(s) involved. We hypothesized that T2D patients would exhibit an augmented muscle metaboreflex activation and this contributes to greater pressor and sympathetic responses during exercise. Mean arterial pressure (MAP), heart rate (HR), and muscle sympathetic nerve activity (MSNA) were measured in 16 patients with T2D (8 normotensive and 8 hypertensive) and 10 healthy controls. Graded isolation of the muscle metaboreflex was achieved by post-exercise ischemia (PEI) following static handgrip performed at 30% and 40% maximal voluntary contraction (MVC). A cold pressor test (CPT) was also performed as a generalized sympatho-excitatory stimulus. Increases in MAP and MSNA during 30 and 40% MVC handgrip were augmented in T2D patients compared to controls ( $P<0.05$ ), and these differences were maintained during PEI (MAP: 30% PEI: T2D,  $\Delta 16 \pm 2$  vs. Controls,  $\Delta 8 \pm 1$  mmHg; 40% PEI: T2D,  $\Delta 26 \pm 3$  vs. Controls,  $\Delta 16 \pm 2$  mmHg, both  $P<0.05$ ). MAP and MSNA responses to handgrip and PEI were not different between normotensive and hypertensive T2D patients ( $P>0.05$ ). Interestingly, MSNA responses were also greater in T2D patients compared to controls during the CPT ( $P<0.05$ ). Collectively, these findings indicate that muscle metaboreflex activation is augmented in T2D patients and this contributes, in part, to augmented pressor and sympathetic responses to exercise in this patient group. Greater CPT responses suggest that a heightened central sympathetic reactivity may be involved.

46 **NEW & NOTEWORTHY**

47       Muscle metaboreflex activation is augmented in type 2 diabetic patients, and this  
48       contributes, in part, to augmented pressor and sympathetic responses to exercise in this patient  
49       group. These findings provide important insight to the neural mechanisms that contribute to the  
50       exaggerated increases in exercise blood pressure in type 2 diabetes.

51

52

## INTRODUCTION

Type 2 diabetes patients (T2D) exhibit exaggerated increases in arterial blood pressure (BP) during exercise (23, 25, 39, 44). Augmented BP responses have been observed even during moderate intensity handgrip (38), a level of isometric forearm muscle contraction that is equivalent to many activities of daily living such as opening jars, or carrying groceries. This is important because repeated surges in BP throughout the day have been related to increased cardiovascular risk (11, 37). Likewise, exaggerated increases in exercise BP are related to adverse cardiovascular and cerebrovascular events both during and after physical activity (19, 27, 33). Indeed, the incidence of cardiovascular and cerebrovascular events such as myocardial infarction and stroke are significantly elevated among T2D patients (5, 26, 28, 57). An augmented pressor response to exercise is also a predictor for the development of hypertension (HTN) (10, 47), a common comorbidity among T2D patients (1, 3, 49, 50). However, despite exaggerated BP responses to exercise and the associated increase in morbidity and mortality in T2D, little is known regarding the underlying neural mechanism(s) involved.

Exercise evokes increases in BP, muscle sympathetic nerve activity (MSNA), and heart rate (HR) that are a result of an integration of central signals originating from higher brain centers (i.e., central command) (15), feedback signals from mechanically and metabolically sensitive afferents in contracting skeletal muscle (i.e., exercise pressor reflex; EPR) (2), and input from the arterial and cardiopulmonary baroreceptors (14, 21). During static handgrip exercise, central command increases heart rate and cardiac output by withdrawing parasympathetic tone (15), whereas the metabolic component of the EPR (i.e., muscle metaboreflex) is primarily responsible for the intensity-dependent increase in MSNA and peripheral vasoconstriction (30, 45). A number of studies have examined the muscle

metaboreflex in subjects with risk factors for T2D (e.g., obesity) and have yielded mixed results (9, 29, 35, 40, 41, 52). Surprisingly, there is a paucity of studies examining the muscle metaboreflex in subjects with overt T2D. Furthermore, no studies have examined whether muscle metaboreflex activation in T2D leads to excessive MSNA responses that could contribute to an exaggerated EPR. A focus on the regulatory mechanisms underlying the augmented neural cardiovascular responses to exercise in T2D is important and clinically relevant.

Given the vital contribution of the muscle metaboreflex to the BP response to exercise, and the previous work demonstrating exaggerated BP responses to exercise in T2D patients, the purpose of this study was to test the hypothesis that BP and MSNA responses to muscle metaboreflex activation would be greater in T2D patients compared to healthy control subjects. Additionally, because T2D is commonly associated with HTN, and previous work has shown that the muscle metaboreflex is augmented with HTN (9, 34, 41), we also hypothesized that muscle metaboreflex activation would be further enhanced in T2D patients with HTN. To test these hypotheses, BP, MSNA, and HR were measured during graded isolation of the muscle metaboreflex using post-exercise ischemia (PEI) following static handgrip performed at 30% and 40% maximal voluntary contraction (MVC). PEI was used to trap local metabolites produced during exercise and isolate activation of metabolically sensitive skeletal muscle afferent nerve endings from the mechanical component of the EPR and central command (2, 30). A cold pressor test (CPT) was also performed to quantify BP and MSNA responses to a generalized non-exercise sympatho-excitatory stimulus.

## Methods

*Subjects.* A total of 27 subjects participated in the present study: Sixteen patients with T2D (reported duration of disease:  $8 \pm 2$  years) and 10 healthy controls matched to T2D patients for age, sex and body weight. General baseline characteristics of the T2D patients and healthy control subjects are provided in Table 1. Eight of the T2D patients also had a clinical diagnosis of hypertension. These patients were all being treated for their hypertension but we excluded any patients taking medications directly influencing MSNA (e.g., central sympathoinhibitors such as clonidine). A listing of the medications being taken by the T2D patients is provided in Table 2. Importantly, none of the T2D patients were being treated for or had symptoms of peripheral neuropathy. Table 3 provides a comparison of baseline characteristics between the T2D patients with and without hypertension. Each subject received a verbal and written explanation of the goals of the study, the experimental measurements, and risks and benefits associated with the study after which each subject provided written informed consent. All subjects also completed a medical health history questionnaire and a 12-h fasting blood chemistry screening including a lipid panel and a metabolic panel that also includes insulin, glucose, and Hb<sub>A1c</sub> measurement. The experimental procedures and protocols used conformed to the Declaration of Helsinki and were approved by the University of Missouri Health Sciences Institutional Review Board.

*Cardiovascular and Metabolic Measurements.* HR and BP were continuously monitored using a lead II surface ECG (Q710; Quinton, Bothell, WA, USA) and a servo-controlled finger photoplethysmography (Finometer; Finapres Medical Systems, Amsterdam, The Netherlands), respectively. For Finometer measurements, return to flow calibrations were performed and

physiocal turned off before each recording. The changes in BP measured using the Finometer have been shown to provide an accurate estimate of directly measured intra-arterial BP (18, 46). Also, an automated sphygmomanometer (Welch Allyn, Skaneateles Falls, NY) recorded resting BP by the auscultation of the brachial artery of the right arm for absolute values of BP and to validate BP measurements from the Finometer (7, 54). Respiratory movements were monitored using a strain-guage pneumograph placed around the abdomen (Pneumotrace, UFI, Morro Bay, CA, USA) to avoid potential confound of large respiratory excursions on cardiovascular measurements during handgrip and PEI. Insulin was measured via an EIA assay (ALPCO, Salem, NH). Insulin resistance was assessed using the homeostatic model assessment of insulin resistance (HOMA-IR):  $\text{HOMA-IR} = (\text{glucose} \times \text{insulin}) / 22.5$ .

*Muscle Sympathetic Nerve Activity.* Multiunit postganglionic MSNA was recorded using standard microneurographic techniques, as previously described (13, 36, 53, 54). Briefly, a tungsten microelectrode was placed into the peroneal nerve near the left fibular head, and a reference microelectrode was inserted 2-3 cm away. Signals were amplified, filtered (bandwidth 0.7-2.0 kHz), rectified and integrated (0.1 s time constant) to obtain mean voltage neurograms using a nerve traffic analyzer (Nerve Traffic analyzer, model 662c-3; University of Iowa Bioengineering, Iowa City, IA). MSNA was identified by the presence of spontaneous pulse synchronous bursts that were responsive to end-expiratory breath holds, but not to arousal or stroking of the skin. Although MSNA signals were obtained in all control subjects, 1 normotensive and 1 hypertensive T2D patient were highly sensitive to the procedure so it was stopped, and we were unable to attain quality signals in 2 others (1 normotensive and 1



hypertensive T2D patient). All neural cardiovascular data was acquired at a frequency of 1,000 Hz using Chart version 5.2 (Powerlab, ADInstruments, Bella Vista, NSW, Australia).

*Isometric handgrip.* Subjects were seated in a semi-recumbent position with a handgrip dynamometer held in the right hand (model 76618; Lafayette Instrument, Lafayette, IN) with the limb supported on an adjustable bedside table. Maximum voluntary contraction (MVC) was determined as the highest of three to five maximal efforts each separated by 1 min, and was used to calculate relative work rates of 30 and 40% MVC for the experimental protocol. During the experimental protocol, ratings of perceived exertion (RPE) were acquired using the Borg scale of 6 to 20 at the end of each bout of handgrip.

*Experimental protocol.* All experiments were performed in a dimly-lit room at an ambient room temperature of 22-24°C with external stimuli minimized. On the experimental day, subjects arrived at the laboratory following an overnight fast, and were also requested to abstain from caffeinated beverages for 12 h and strenuous physical activity and alcohol for at least 24 h. T2D patients were also instructed to refrain from all medication use the morning of the study. Before the performance of the experimental protocol, each subject was familiarized with all measurements, the equipment and testing procedures.

After instrumentation for all experimental measurements, a 10 min baseline recording was performed to determine resting cardiovascular variables and MSNA. Subjects then performed 2 min of isometric handgrip at either 30% or 40% MVC followed by 2 min and 15 s of forearm ischemia to isolate muscle metaboreflex activation (PEI). PEI was achieved by inflation of a blood pressure cuff around the upper arm to suprasystolic pressure ( $>240$  mmHg) 5

s before the end of handgrip exercise. The additional 15 s of PEI was included to account for the initial decrease in BP and MSNA that occurs immediately following the cessation of handgrip exercise. Visual feedback regarding the handgrip force exerted was provided via a personal computer displayed at eye level (Chart v5.2, Powerlab). In all cases except one, the 30% MVC trial was performed first due to the greater probability for muscle tension and loss of the MSNA signal during 40% MVC handgrip. The handgrip trials were separated by at least 15 min to allow BP, MSNA, and HR to return to baseline values.

*Cold pressor test.* A cold pressor test was used to determine BP, MSNA, and HR responses to a generalized, non-exercise sympathoexcitatory stimulus (55). The right hand was placed in ice water for 2 min. All variables were recorded during a 2 min baseline period, during the cold pressor test, and for 2 min during recovery.

*Data Analysis.* Resting values for BP, MSNA, and HR were calculated as mean values over a 10 min steady-state period. MSNA was analyzed using a custom LabVIEW program (12, 13). MSNA was quantified as burst frequency (bursts/min), burst incidence (bursts/100 cardiac cycles) and total activity (burst frequency multiplied by mean burst amplitude; AU/min). To account for variation in burst amplitude, MSNA burst amplitudes were expressed as a percentage of the average of the three largest bursts during baseline (assigned a mean value of 100 arbitrary units; AU). Thirty second averages of handgrip exercise (30-60 s and 90-120 s) and the final 60 s averages of PEI were used for group comparisons. The first 60 s and second 60 s of the cold pressor test were averaged and used for group comparisons.

To examine the interaction between the muscle metaboreflex and the arterial baroreflex, spontaneous baroreflex control of MSNA was calculated during PEI and compared to resting measures. Briefly, MSNA was averaged over 3-mmHg diastolic BP ranges (bins), and a weighted linear regression analysis between the spontaneous changes in MSNA and diastolic BP was performed. MSNA within each pressure bin was calculated as total MSNA (total area of all MSNA bursts relative to the number of cardiac cycles) and expressed as AU/beat. Burst incidence within each pressure bin was also calculated. Diastolic BP was used for this analysis because changes in MSNA correlate closely with changes in diastolic BP but not systolic BP (51).

*Statistical Analysis.* All data are reported as mean  $\pm$  SEM. Statistical comparisons of resting physiological variables between groups were made using one-way analysis of variance (ANOVA). Statistical comparisons of changes in BP, MSNA, and HR between groups during handgrip and PEI, and during the CPT, were made using two-way repeated measures ANOVA. Bonferonni post hoc testing was applied where significant main effects were found. Pearson product-moment correlation coefficients were performed between metabolic parameters and BP and MSNA responses to handgrip, PEI and the CPT. Data was analyzed using SigmaPlot 13 (Systat Software Inc.).

## **Results**

*Subject Characteristics.* Age and BMI were not different between Controls and T2D patients (Table 1). As expected, T2D patients had significantly elevated plasma glucose, HbA1c, and

HOMA-IR compared to Control subjects (Table 1). No significant differences in resting systolic, diastolic, or MAP were found between Controls and T2D Patients (Table 1). In this regard, all hypertensive T2D patients were currently on an active treatment regimen ( $\geq 1$  antihypertensive medications) (Table 2). Resting MSNA burst frequency and burst incidence was also not different between Controls and T2D Patients (Table 1). A comparison of normotensive and hypertensive T2D patients demonstrated no significant differences in resting metabolic, cardiovascular or MSNA variables (Table 3). The only significant difference found was a greater BMI in hypertensive T2D patients. MVC was not different between groups (Control:  $40 \pm 3$  kg; T2D:  $38 \pm 3$  kg; T2D+HTN:  $42 \pm 3$  kg;  $P=0.908$ ).

#### *Isometric handgrip and PEI.*

Original recordings of BP and MSNA at baseline, during 30% MVC handgrip, and during PEI in 3 T2D patients and 3 control subjects are displayed in Fig. 1. The increase in MAP was significantly greater during 30% and 40% MVC handgrip in T2D patients compared to control subjects and these augmented pressor responses in T2D patients were maintained during PEI (Fig. 2). Similar results were found with systolic BP (30% MVC,  $P=0.001$  vs. Control; 40% MVC,  $P=0.011$  vs. Control), and diastolic BP (30% MVC,  $P<0.001$  vs. Control; 40% MVC,  $P=0.021$  vs. Control) (data not shown). The increase in HR during handgrip was also significantly greater in T2D patients, but only during 30% MVC handgrip ( $P<0.001$  vs. Control), and returned toward baseline values during PEI following both 30% and 40% handgrip in both T2D patients and controls (Figure 2). RPE values obtained at the end of handgrip were not different between groups (30% MVC: T2D,  $13.5 \pm 0.6$  vs. Control,  $11.8 \pm 0.6$ ,  $P=0.057$ ; 40% MVC: T2D,  $15.7 \pm 0.5$  vs. Control,  $14.3 \pm 0.8$ ,  $P=0.161$ ).

MSNA responses to handgrip and PEI were also significantly greater in T2D patients compared to controls (Fig. 3). In this regard, during handgrip exercise at 30% MVC, the change in MSNA burst frequency and percent change in total activity was augmented in T2D patients compared to controls, and this augmented MSNA response was sustained during PEI (Fig 3A). Likewise, MSNA burst incidence was greater in T2D patients during 30% MVC handgrip and PEI (HG 120: T2D,  $\Delta 20.1 \pm 3.8$  vs. Control,  $\Delta 6.5 \pm 1.7$  burst/100 heartbeats,  $P < 0.001$ ; PEI: T2D,  $\Delta 23.8 \pm 6.5$  vs. Control,  $\Delta 9.9 \pm 2.5$  burst/100 heartbeats,  $P < 0.001$ ). During handgrip and PEI at 40% MVC, the percent change in MSNA total activity was also augmented in T2D patients; whereas, the change in MSNA burst frequency did not reach statistical significance, although there was a tendency for a greater response in T2D patients (Fig. 3B, top panel). The latter may be due to maintaining MSNA recordings in only 8 T2D patients during 40% MVC handgrip. This was primarily due to muscle tension and loss of the MSNA signal with this higher intensity of handgrip. In contrast, quality MSNA recordings were maintained during 30% MVC handgrip and PEI in 12 T2D patients. Nevertheless, MSNA burst incidence was greater in the T2D patients during 40% MVC handgrip and PEI (HG 120: T2D,  $\Delta 27.4 \pm 6.1$  vs. Control,  $\Delta 13.0 \pm 5.6$  burst/100 heartbeats; PEI: T2D,  $\Delta 31.7 \pm 4.9$  vs. Control,  $\Delta 16.7 \pm 3.6$  burst/100 heartbeats,  $P = 0.04$ ).

Among the T2D patients, BP and MSNA responses to isometric handgrip at 30% MVC were similar between those with and without hypertension. Likewise, BP responses to 40% MVC handgrip were not different between normotensive and hypertensive T2D patients, whereas the MSNA response to 40% MVC handgrip was greater in hypertensive T2D patients. Nevertheless, BP and MSNA responses during PEI following both 30% and 40% MVC were not different between normotensive and hypertensive T2D patients (Fig 4). We also tested for

potential sex differences in cardiovascular responses to handgrip and PEI, since our groups were composed of both men and women. We found no effect of sex on any of the variables of interest both during handgrip and PEI. For example, in the control group (N=5 men and 5 women), the increase in MSNA during PEI following 30% MVC handgrip was  $\Delta 5.2 \pm 1.5$  bursts/min in the men and  $\Delta 6.4 \pm 2.6$  bursts/min in the women ( $P=0.702$ ) and in the T2D patients (N=6 men and 6 women), the increase in MSNA during PEI was  $\Delta 17 \pm 5.4$  bursts/min in the men and  $\Delta 14.5 \pm 3.1$  burst/min in the women ( $P=0.615$ ).

The increases in MSNA total activity during PEI following both 30% MVC and 40% MVC handgrip were significantly correlated with fasting glucose, HbA1c, and HOMA-IR (Figure 5). In contrast, weaker relationships between fasting insulin and MSNA responses during PEI were found (30% MVC PEI:  $R=0.2$ ,  $P=0.417$ ; 40% MVC PEI:  $R=0.4$ ,  $P=0.153$ ).

Spontaneous baroreflex control of MSNA at rest was not different between T2D patients and controls for either burst incidence or total MSNA (burst incidence: T2D,  $-4.7 \pm 0.4$  vs. Control,  $-4.3 \pm 0.5$  bursts $\cdot 100\text{hb}^{-1}\cdot\text{mmHg}^{-1}$ ,  $P=0.534$ ; total MSNA: T2D,  $-2.5 \pm 0.2$  vs. Control,  $-2.5 \pm 0.3$  AU $\cdot\text{beat}^{-1}\cdot\text{mmHg}^{-1}$ ,  $P=0.936$ ). Likewise, the increase in total MSNA gain during PEI was not different between groups (30% MVC PEI: T2D,  $-3.4 \pm 0.6$  vs. Control,  $-4.4 \pm 0.3$  AU $\cdot\text{beat}^{-1}\cdot\text{mmHg}^{-1}$ ,  $P=0.323$ ).

*Cold pressor test.* Although the increase in MAP during the cold pressor test appeared to be greater in T2D patients (n=9) compared to control subjects (n=9), this did not reach statistical significance (Fig. 6). However, the change in MSNA burst frequency (Fig 6B) and MSNA total activity (CPT min 2: T2D,  $151 \pm 20$  vs. Control,  $55 \pm 21$  %AU/min,  $P=0.005$ ) were significantly greater in T2D patients (n=7) compared to control subjects (n=9). HR responses to the cold

pressor test were not different between groups (CPT min 2: T2D,  $\Delta 6 \pm 3$  vs. Control,  $\Delta 3 \pm 2$  bpm,  $P=0.330$ ).

For the CPT, significant correlations were noted between increases in MSNA and fasting glucose ( $R=0.55$ ,  $P=0.027$ ) and HOMA-IR ( $R=0.79$ ,  $P=0.001$ ), but not HbA1c ( $R=0.39$ ,  $P=0.134$ ) or fasting insulin ( $R=0.35$ ,  $P=0.218$ ). Interestingly, no significant correlations were observed between the increases in MSNA during the CPT and during PEI following 30% MVC ( $R=0.368$ ,  $P=0.161$ ) or 40% MVC ( $R=0.472$ ,  $P=0.103$ ).

## Discussion

The major and novel finding of the present study is that T2D patients exhibit a heightened activation of the metabolic component of the EPR. Indeed, augmented pressor and MSNA responses during handgrip were maintained during isolation of the muscle metaboreflex with PEI. Thus, greater MSNA and BP responses remained in T2D patients when input from central command and the muscle mechanoreflex were removed. Notably, MSNA responses were also greater in T2D patients compared to controls during the CPT. Collectively, these findings indicate, for the first time, that the metabolic component of the EPR is augmented with T2D and this contributes, in part, to augmented pressor and sympathetic responses to exercise in T2D patients. Greater MSNA responses to a generalized non-exercise sympatho-excitatory stimulus such as the CPT suggest that a heightened central sympathetic reactivity may be involved.

Given the fairly well-documented augmentation in exercise BP in T2D patients (23, 25, 38, 39, 43, 44), it was surprising that few studies have attempted to examine potential alterations in the underlying neural cardiovascular mechanisms in this patient group. Furthermore, to our knowledge, no studies have measured MSNA responses during exercise in T2D patients. Given

the significance of the muscle metaboreflex to the pressor response to exercise, we chose to begin with the muscle metaboreflex. To this end, graded PEI was used to trap local metabolites produced by active skeletal muscle and preserve activation of metabosensitive afferent nerve endings, and therefore isolate the metabolic component of the EPR (2, 30). We now demonstrate that PEI following 30 and 40% MVC handgrip resulted in augmented pressor responses, and that these augmented pressor responses were accompanied by enhanced increases in MSNA. Interestingly, the increase in MSNA during PEI was significantly correlated with glucose control and insulin resistance markers (i.e., fasting glucose, HbA1c, and HOMA-IR), implying that the effectiveness of T2D control may play a role (see Figure 5). Indeed, the higher the fasting glucose and HbA1c and the greater level of insulin resistance, the greater augmentation in muscle metaboreflex activation. Taken together, our results suggest that the effects of T2D on the regulatory mechanisms underlying the neural cardiovascular responses to exercise involve at least a heightened metabolic component of the EPR that appears to be related to the severity of T2D.

T2D is commonly associated with hypertension (1, 3, 49, 50), and indeed, a significant number of T2D patients recruited for the present study had hypertension. Given that both human and animal studies have suggested an exaggerated activation of the muscle metaboreflex in hypertension (9, 34, 40, 41), we compared responses between normotensive and hypertensive T2D patients to test if the co-existence of hypertension and T2D would further augment the pressor and MSNA responses to PEI. However, we found that the heightened metaboreflex activation observed in the T2D patients was unaffected by hypertensive status (see figure 4). Both the BP and MSNA responses to handgrip and PEI were similar between normotensive and hypertensive T2D patients. Nonetheless, it is important to note that our data are only reflective of



hypertensive T2D patients that have well controlled BP and it is possible that uncontrolled hypertensive T2D patients or never treated hypertensive T2D patients might have different responses. Additional studies are warranted in this regard to further understand the influence of uncontrolled hypertension among T2D patients on muscle metaboreflex activation.

The mechanisms responsible for the exaggerated muscle metaboreflex activation in T2D are not entirely clear. Although elevations in BP and MSNA during muscle metaboreflex activation are primarily driven by the muscle metaboreflex, there is an interaction between the metaboreflex and the arterial baroreflex that can modify such responses. Indeed, studies have shown exaggerated neural cardiovascular responses when input from the arterial baroreflex is removed (48, 56). Likewise, in healthy humans, an increase in the baroreflex control of MSNA has been observed during isolation of the muscle metaboreflex with PEI (8, 20, 22). Since previous work suggests that the sensitivity of the arterial baroreflex may be impaired in conditions associated with T2D such as obesity (6, 16) and hypertension (17, 31), we examined the interaction between the muscle metaboreflex and the arterial baroreflex. Our findings suggest that an impaired baroreflex control of MSNA does not appear to contribute to the augmented MSNA and BP responses observed during PEI since both groups exhibited an increase in MSNA baroreflex sensitivity with PEI, similar to previous studies (8, 20, 22). However, since only spontaneous baroreflex measures were used, additional studies are needed to more fully characterize arterial baroreflex function. It also remains possible that the group IV afferent fibers in the skeletal muscle interstitium that are responsive to changes in metabolic concentrations have greater sensitivity in T2D. Alternatively, although handgrip strength and perceived exertion were not different between T2D patients and control subjects, it remains possible that T2D patients experience greater metabolite build-up within the muscle interstitium during handgrip.

In this regard, previous work suggests altered skeletal muscle metabolism in T2D patients (4, 42), which may lead to greater production of substances during muscle contraction that stimulate skeletal muscle afferents and contribute to greater metaboreflex activation in T2D patients. Identification of the particular substances responsible for stimulating muscle afferent remains an ongoing area of research (24), and future studies would be needed to characterize the responsiveness of skeletal muscle afferents to various substances in T2D, likely including animal investigations.

In the present study, a CPT was used as a generalized sympathoexcitatory stimulus to assess whether a heightened central sympathetic activation may be augmented in T2D patients. Interestingly, the MSNA responses to the CPT were greater in T2D patients, and there was also a trend for a greater pressor response but this did not reach statistical significance. These results suggest that heightened sympathetic responsiveness in T2D may be global and not specific to metaboreflex activation. However, when MSNA responses to the CPT were compared to MSNA responses to isolated metaboreflex activation, no significant correlations were found. Also, correlations between metabolic measures (fasting glucose, HbA1c, and HOMA-IR) and the MSNA responses to the cold pressor test were noticeably weaker than was seen with isolated metaboreflex activation. Although not determining causality or lack thereof, these data suggest that greater central sympathetic activation may not completely account for the augmented metaboreflex activation of MSNA observed in T2D patients. Nevertheless, further studies are warranted to investigate the mechanism(s) responsible for the heightened BP and MSNA mediated metaboreflex responses in T2D as well as hypertension and will likely require animal investigations to tease apart the afferent, central and efferent pathways.

## Perspectives

Exaggerated increases in exercise BP are related to adverse cardiovascular and cerebrovascular events both during and following exercise (19, 27, 33). Although it is known that T2D patients exhibit augmented BP responses to exercise, limited studies have focused on the sympathetic and cardiovascular responses to isometric exercise in this patient group. This is important because isometric contractions are a component of many daily activities, and are capable of inducing marked increases in BP and afterload on the heart even when performed with a small muscle mass (32). This highlights the significance of the augmented increases in BP and MSNA that were observed in T2D patients and attributable in part to a heightened muscle metaboreflex activation. These findings are of vital importance given the incidence of myocardial infarction and stroke among T2D patients (5, 26, 28, 57), and the number of common daily activities that involve an isometric muscle contraction component. Given our findings of a significant contribution of the muscle metaboreflex to greater pressor and sympathetic responses to isometric contractions in patients with T2D, future studies to identify the mechanism(s) responsible to target and reduce such hyper-responses are needed. In the meantime, if prescribed to a T2D patient for better health and fitness, resistance exercise training should be prescribed at a low intensity and duration for this patient population.

In summary, we report for the first time that greater pressor responses in T2D during isometric handgrip are attributable, in part, to heightened muscle metaboreflex activation. Augmented pressor responses to handgrip and PEI in T2D patients are paralleled by exaggerated increases in MSNA. These findings demonstrate that the metabolic component of the EPR is augmented in T2D, and provide important insight to the neural mechanisms that contribute to the exaggerated increases in exercise BP in T2D.

## **Grants**

This research was supported by an American Heart Association Grant in Aid (20160072 to P.J.F.), by a Mizzou Advantage Grant (to P.J.F.) and by an ACSM Foundation Research Grant from the American College of Sports Medicine Foundation (S.W.H.). The results of the study do not constitute endorsement by ACSM.

## **Disclosures**

No conflicts of interest, financial or otherwise, are declared by the authors.

## **Author Contributions**

Author contributions: S.W.H., J.P.F., and P.J.F. conception and design of research; C.M., and G.L. assisted with patient recruitment and screening; S.W.H., R.M.R., J.P.F., and P.J.F. performed experiments; S.W.H. analyzed data and prepared figures; S.W.H., J.P.F., and P.J.F. interpreted results of experiments; S.W.H. drafted manuscript; S.W.H., C.M., J.P.F., and P.J.F. edited and revised the manuscript; S.W.H., R.M.R., C.M., G.L., J.P.F., and P.J.F. approved the final version of the manuscript.

## **Acknowledgements**

We thank Charla Jay, R.N., for assistance with screening blood draws and patient recruitment. We also would like to thank Joseph W. LeMaster, M.D., and Colin N. Young, PhD for assistance

417 during the initiation of these studies. This research was submitted in partial fulfilment of the  
418 requirements for the degree of Doctor of Philosophy for Seth W. Holwerda.

419

## References

1. National High Blood Pressure Education Program Working Group report on hypertension in diabetes. *Hypertension*. 1994;23(2):145-58; discussion 59-60.
2. Alam M, Smirk FH. Observations in man upon a blood pressure raising reflex arising from the voluntary muscles. *The Journal of physiology*. 1937;89(4):372-83.
3. Arauz-Pacheco C, Parrott MA, Raskin P. Treatment of hypertension in adults with diabetes. *Diabetes care*. 2003;26 Suppl 1:S80-2.
4. Avogaro A, Toffolo G, Miola M et al. Intracellular lactate- and pyruvate-interconversion rates are increased in muscle tissue of non-insulin-dependent diabetic individuals. *The Journal of clinical investigation*. 1996;98(1):108-15.
5. Barrett-Connor EL, Cohn BA, Wingard DL, Edelstein SL. Why is diabetes mellitus a stronger risk factor for fatal ischemic heart disease in women than in men? The Rancho Bernardo Study. *Jama*. 1991;265(5):627-31.
6. Beske SD, Alvarez GE, Ballard TP, Davy KP. Reduced cardiovagal baroreflex gain in visceral obesity: implications for the metabolic syndrome. *American journal of physiology. Heart and circulatory physiology*. 2002;282(2):H630-5.
7. Credeur DP, Holwerda SW, Boyle LJ, Vianna LC, Jensen AK, Fadel PJ. Effect of Aging on Carotid Baroreflex Control of Blood Pressure and Leg Vascular Conductance in Women. *American journal of physiology. Heart and circulatory physiology*. 2014.
8. Cui J, Wilson TE, Shibasaki M, Hodges NA, Crandall CG. Baroreflex modulation of muscle sympathetic nerve activity during posthandgrip muscle ischemia in humans. *J Appl Physiol (1985)*. 2001;91(4):1679-86.
9. Delaney EP, Greaney JL, Edwards DG, Rose WC, Fadel PJ, Farquhar WB. Exaggerated sympathetic and pressor responses to handgrip exercise in older hypertensive humans: role of the muscle metaboreflex. *American journal of physiology. Heart and circulatory physiology*. 2010;299(5):H1318-27.
10. Dlin RA, Hanne N, Silverberg DS, Bar-Or O. Follow-up of normotensive men with exaggerated blood pressure response to exercise. *American heart journal*. 1983;106(2):316-20.
11. Eguchi K, Hoshida S, Schwartz JE, Shimada K, Kario K. Visit-to-visit and ambulatory blood pressure variability as predictors of incident cardiovascular events in patients with hypertension. *American journal of hypertension*. 2012;25(9):962-8.
12. Fairfax ST, Holwerda SW, Credeur DP et al. The role of alpha-adrenergic receptors in mediating beat-by-beat sympathetic vascular transduction in the forearm of resting man. *The Journal of physiology*. 2013;591(Pt 14):3637-49.
13. Fairfax ST, Padilla J, Vianna LC, Davis MJ, Fadel PJ. Spontaneous bursts of muscle sympathetic nerve activity decrease leg vascular conductance in resting humans. *American journal of physiology. Heart and circulatory physiology*. 2013;304(5):H759-66.
14. Fisher JP, Young CN, Fadel PJ. Autonomic adjustments to exercise in humans. *Comprehensive Physiology*. 2015;5(2):475-512.
15. Goodwin GM, McCloskey DI, Mitchell JH. Cardiovascular and respiratory responses to changes in central command during isometric exercise at constant muscle tension. *The Journal of physiology*. 1972;226(1):173-90.
16. Grassi G, Seravalle G, Dell'Oro R, Turri C, Bolla GB, Mancina G. Adrenergic and reflex abnormalities in obesity-related hypertension. *Hypertension*. 2000;36(4):538-42.
17. Gribbin B, Pickering TG, Sleight P, Peto R. Effect of age and high blood pressure on baroreflex sensitivity in man. *Circulation research*. 1971;29(4):424-31.

18. Harms MP, Wesseling KH, Pott F et al. Continuous stroke volume monitoring by modelling flow from non-invasive measurement of arterial pressure in humans under orthostatic stress. *Clin Sci (Lond)*. 1999;97(3):291-301.
19. Hoberg E, Schuler G, Kunze B et al. Silent myocardial ischemia as a potential link between lack of premonitoring symptoms and increased risk of cardiac arrest during physical stress. *The American journal of cardiology*. 1990;65(9):583-9.
20. Ichinose M, Saito M, Wada H, Kitano A, Kondo N, Nishiyasu T. Modulation of arterial baroreflex control of muscle sympathetic nerve activity by muscle metaboreflex in humans. *American journal of physiology. Heart and circulatory physiology*. 2004;286(2):H701-7.
21. Joyner MJ. Baroreceptor function during exercise: resetting the record. *Experimental physiology*. 2006;91(1):27-36.
22. Kamiya A, Michikami D, Fu Q et al. Static handgrip exercise modifies arterial baroreflex control of vascular sympathetic outflow in humans. *American journal of physiology. Regulatory, integrative and comparative physiology*. 2001;281(4):R1134-9.
23. Karavelioglu Y, Karapinar H, Gul I et al. Blood pressure response to exercise is exaggerated in normotensive diabetic patients. *Blood pressure*. 2013;22(1):21-6.
24. Kaufman MP, Hayes SG. The exercise pressor reflex. *Clinical autonomic research : official journal of the Clinical Autonomic Research Society*. 2002;12(6):429-39.
25. Kingwell BA, Formosa M, Muhlmann M, Bradley SJ, McConell GK. Type 2 diabetic individuals have impaired leg blood flow responses to exercise: role of endothelium-dependent vasodilation. *Diabetes care*. 2003;26(3):899-904.
26. Kissela BM, Khoury J, Kleindorfer D et al. Epidemiology of ischemic stroke in patients with diabetes: the greater Cincinnati/Northern Kentucky Stroke Study. *Diabetes care*. 2005;28(2):355-9.
27. Kokkinos PF, Andreas PE, Coutoulakis E et al. Determinants of exercise blood pressure response in normotensive and hypertensive women: role of cardiorespiratory fitness. *Journal of cardiopulmonary rehabilitation*. 2002;22(3):178-83.
28. Koskinen P, Manttari M, Manninen V, Huttunen JK, Heinonen OP, Frick MH. Coronary heart disease incidence in NIDDM patients in the Helsinki Heart Study. *Diabetes care*. 1992;15(7):820-5.
29. Limberg J, Morgan B, Schrage W. Mechanical and metabolic reflex activation of the sympathetic nervous system in younger adults with metabolic syndrome. *Autonomic neuroscience : basic & clinical*. 2014;183:100-5.
30. Mark AL, Victor RG, Nerhed C, Wallin BG. Microneurographic studies of the mechanisms of sympathetic nerve responses to static exercise in humans. *Circulation research*. 1985;57(3):461-9.
31. Matsukawa T, Gotoh E, Hasegawa O, Shionoiri H, Tochikubo O, Ishii M. Reduced baroreflex changes in muscle sympathetic nerve activity during blood pressure elevation in essential hypertension. *Journal of hypertension*. 1991;9(6):537-42.
32. Mitchell JH, Wildenthal K. Static (isometric) exercise and the heart: physiological and clinical considerations. *Annual review of medicine*. 1974;25:369-81.
33. Mittleman MA, Siscovick DS. Physical exertion as a trigger of myocardial infarction and sudden cardiac death. *Cardiology clinics*. 1996;14(2):263-70.
34. Mizuno M, Murphy MN, Mitchell JH, Smith SA. Skeletal muscle reflex-mediated changes in sympathetic nerve activity are abnormal in spontaneously hypertensive rats. *American journal of physiology. Heart and circulatory physiology*. 2011;300(3):H968-77.

- 512 35. Negrao CE, Trombetta IC, Batalha LT et al. Muscle metaboreflex control is diminished in  
513 normotensive obese women. *American journal of physiology. Heart and circulatory physiology.*  
514 2001;281(2):H469-75.
- 515 36. Padilla J, Young CN, Simmons GH et al. Increased muscle sympathetic nerve activity acutely  
516 alters conduit artery shear rate patterns. *American journal of physiology. Heart and circulatory*  
517 *physiology.* 2010;298(4):H1128-35.
- 518 37. Parati G, Ochoa JE, Bilo G. Blood pressure variability, cardiovascular risk, and risk for renal  
519 disease progression. *Current hypertension reports.* 2012;14(5):421-31.
- 520 38. Petrofsky JS, Stewart B, Patterson C, Cole M, Al Maly A, Lee S. Cardiovascular responses and  
521 endurance during isometric exercise in patients with Type 2 diabetes compared to control  
522 subjects. *Medical science monitor : international medical journal of experimental and clinical*  
523 *research.* 2005;11(10):CR470-7.
- 524 39. Regensteiner JG, Bauer TA, Reusch JE et al. Cardiac dysfunction during exercise in uncomplicated  
525 type 2 diabetes. *Medicine and science in sports and exercise.* 2009;41(5):977-84.
- 526 40. Rondon MU, Laterza MC, de Matos LD et al. Abnormal muscle metaboreflex control of  
527 sympathetic activity in never-treated hypertensive subjects. *American journal of hypertension.*  
528 2006;19(9):951-7.
- 529 41. Sausen MT, Delaney EP, Stillabower ME, Farquhar WB. Enhanced metaboreflex sensitivity in  
530 hypertensive humans. *European journal of applied physiology.* 2009;105(3):351-6.
- 531 42. Scheuermann-Freestone M, Madsen PL, Manners D et al. Abnormal cardiac and skeletal muscle  
532 energy metabolism in patients with type 2 diabetes. *Circulation.* 2003;107(24):3040-6.
- 533 43. Schultz MG, Hordern MD, Leano R, Coombes JS, Marwick TH, Sharman JE. Lifestyle change  
534 diminishes a hypertensive response to exercise in type 2 diabetes. *Medicine and science in*  
535 *sports and exercise.* 2011;43(5):764-9.
- 536 44. Scott JA, Coombes JS, Prins JB, Leano RL, Marwick TH, Sharman JE. Patients with type 2 diabetes  
537 have exaggerated brachial and central exercise blood pressure: relation to left ventricular  
538 relative wall thickness. *American journal of hypertension.* 2008;21(6):715-21.
- 539 45. Seals DR. Sympathetic neural discharge and vascular resistance during exercise in humans. *J*  
540 *Appl Physiol (1985).* 1989;66(5):2472-8.
- 541 46. Shi X, Gallagher KM, SA SM, Bryant KH, Raven PB. Diminished forearm vasomotor response to  
542 central hypervolemic loading in aerobically fit individuals. *Medicine and science in sports and*  
543 *exercise.* 1996;28(11):1388-95.
- 544 47. Singh JP, Larson MG, Manolio TA et al. Blood pressure response during treadmill testing as a risk  
545 factor for new-onset hypertension. The Framingham heart study. *Circulation.* 1999;99(14):1831-  
546 6.
- 547 48. Smith SA, Williams MA, Leal AK, Mitchell JH, Garry MG. Exercise pressor reflex function is altered  
548 in spontaneously hypertensive rats. *The Journal of physiology.* 2006;577(Pt 3):1009-20.
- 549 49. Sowers JR, Epstein M. Diabetes mellitus and associated hypertension, vascular disease, and  
550 nephropathy. An update. *Hypertension.* 1995;26(6 Pt 1):869-79.
- 551 50. Sowers JR, Epstein M, Frohlich ED. Diabetes, hypertension, and cardiovascular disease: an  
552 update. *Hypertension.* 2001;37(4):1053-9.
- 553 51. Sundlof G, Wallin BG. Human muscle nerve sympathetic activity at rest. Relationship to blood  
554 pressure and age. *The Journal of physiology.* 1978;274:621-37.
- 555 52. Trombetta IC, Batalha LT, Rondon MU et al. Weight loss improves neurovascular and muscle  
556 metaboreflex control in obesity. *American journal of physiology. Heart and circulatory*  
557 *physiology.* 2003;285(3):H974-82.
- 558 53. Vallbo AB, Hagbarth KE, Torebjork HE, Wallin BG. Somatosensory, proprioceptive, and  
559 sympathetic activity in human peripheral nerves. *Physiological reviews.* 1979;59(4):919-57.



54. Vianna LC, Hart EC, Fairfax ST, Charkoudian N, Joyner MJ, Fadel PJ. Influence of age and sex on the pressor response following a spontaneous burst of muscle sympathetic nerve activity. *American journal of physiology. Heart and circulatory physiology*. 2012;302(11):H2419-27.
55. Victor RG, Leimbach WN, Jr., Seals DR, Wallin BG, Mark AL. Effects of the cold pressor test on muscle sympathetic nerve activity in humans. *Hypertension*. 1987;9(5):429-36.
56. Waldrop TG, Mitchell JH. Effects of barodenervation on cardiovascular responses to static muscular contraction. *The American journal of physiology*. 1985;249(4 Pt 2):H710-4.
57. Zhao W, Katzmarzyk PT, Horswell R, Wang Y, Johnson J, Hu G. Sex differences in the risk of stroke and HbA(1c) among diabetic patients. *Diabetologia*. 2014;57(5):918-26.

Figure 1. Original recordings of muscle sympathetic nerve activity (MSNA) and mean arterial pressure (MAP) in 3 type 2 diabetic patients (T2D; Panels A-C) and 3 control subjects (Panels D-F) at baseline, during 30% MVC isometric handgrip, and during post-exercise ischemia (PEI).

Figure 2. Mean and individual data showing the change in mean arterial pressure (MAP) and heart rate (HR) at 60 and 120 s of 30% MVC (Panel A) and 40% MVC (Panel B) handgrip followed by subsequent periods of post-exercise ischemia (PEI) in type 2 diabetic patients (T2D) and control subjects. \* $P < 0.05$  vs. Control.

Figure 3. Mean and individual data showing the change in muscle sympathetic nerve activity (MSNA) at 60 and 120 s of 30% MVC (Panel A) and 40% MVC (Panel B) handgrip followed by subsequent periods of post-exercise ischemia (PEI) in type 2 diabetic patients (T2D) and control subjects. \* $P < 0.05$  vs. Control.

Figure 4. Mean summary data showing the changes in mean arterial pressure (MAP) and muscle sympathetic nerve activity (MSNA) during isolation of the muscle metaboreflex with post-exercise ischemia (PEI) following 30% maximal voluntary contraction (MVC) handgrip (Panel A) and 40% MVC handgrip (Panel B) in normotensive (T2D+NTN) and hypertensive type 2 diabetic patients (T2D+HTN).

Figure 5. Correlations between the change in muscle sympathetic nerve activity (MSNA) during post-exercise ischemia (PEI) following 30% maximal voluntary contraction (MVC) handgrip (Panel A) and 40% MVC handgrip (Panel B) and fasting glucose, glycated hemoglobin (HbA1c), and homeostatic model assessment of insulin resistance (HOMA-IR) in all subjects.

Figure 6. Mean summary data showing the change in mean arterial pressure (MAP; Panel A) and muscle sympathetic nerve activity (MSNA; Panel B) at 60 and 120 s of a cold pressor test in type 2 diabetic patients (T2D) and control subjects. \* $P < 0.05$  vs. Control.

Table 1. Main subject characteristics

	Control	T2D	P value
Sex, men/women	5/5	9/7	
Age, years	46 ± 3	50 ± 2	0.334
BMI, kg/m <sup>2</sup>	29 ± 2	31 ± 4	0.312
Glucose, mg/dL	95 ± 2	198 ± 22*	0.001
HbA1c, %	5.3 ± 0.1	8.6 ± 0.5*	<0.001
Insulin (μIU/mL)	7.6 ± 0.7	11 ± 2.3	0.293
HOMA-IR	1.8 ± 0.2	4.6 ± 0.7*	0.013
Triglycerides, mg/dL	114 ± 20	184 ± 38	0.172
Cardiovascular variables			
Heart rate (bpm)	63 ± 4	69 ± 3	0.212
Systolic BP (mmHg)	122 ± 4	128 ± 4	0.313
Diastolic BP (mmHg)	78 ± 2	81 ± 3	0.394
Mean BP (mmHg)	91 ± 3	97 ± 3	0.258
MSNA			
	N=10	N=12	
Burst frequency (burst/min)	25 ± 4	31 ± 2	0.230
Burst incidence (burst/100hb)	39 ± 6	46 ± 4	0.361
Total activity (AU/min)	1241 ± 219	1411 ± 139	0.520

\*P&lt;0.05 vs. Control

Table 2. Subject medications


	Control	T2D+NTN	T2D+HTN
Hypoglycemic medications			
Biguanide	N=0	N=5	N=7
Sulfonylurea	N=0	N=1	N=0
DPP-4 inhibitor	N=0	N=0	N=3
Insulin	N=0	N=2	N=4
Cardiovascular medications			
ACE inhibitor	N=0	N=0	N=5
Ang II receptor blocker	N=0	N=0	N=1
Diuretic	N=0	N=0	N=4
$\beta$ -blocker	N=0	N=0	N=1
Statin	N=0	N=2	N=5

Table 3. T2D subject characteristics

	T2D+NTN	T2D+HTN	P value
Sex, men/women	4/4	5/3	
Age, years	48 ± 4	51 ± 2	0.576
BMI, kg/m <sup>2</sup>	29 ± 2	34 ± 1†	0.032
Glucose, mg/dL	209 ± 34	187 ± 29	0.637
HbA1c, %	8.8 ± 0.8	8.5 ± 0.7	0.797
Insulin (μIU/mL)	11.4 ± 4.9	10.8 ± 1.5	0.904
HOMA-IR	4.5 ± 1.3	4.8 ± 0.8	0.860
Triglycerides, mg/dL	175 ± 44	92 ± 64	0.833
Cardiovascular variables			
Heart rate (bpm)	69 ± 3	68 ± 4	0.890
Systolic BP (mmHg)	126 ± 6	130 ± 5	0.551
Diastolic BP (mmHg)	80 ± 4	83 ± 5	0.604
Mean BP (mmHg)	95 ± 4	99 ± 5	0.568
MSNA			
	N=6	N=6	
Burst frequency (burst/min)	31 ± 3	31 ± 3	0.980
Burst incidence (burst/100hb)	46 ± 5	45 ± 7	0.955
Total activity (AU/min)	1382 ± 209	1439 ± 201	0.846


†P&lt;0.05 vs. T2D+NTN

Figure 1



**Figure 2**

**A) 30% MVC**



**B) 40% MVC**







Figure 3

A) 30% MVC



B) 40% MVC










Figure 4




**Figure 5**

**A) 30% MVC PEI**



**B) 40% MVC PEI**



**Figure 6**

